



PHARMACY COVERAGE GUIDELINES
SECTION: DRUGS

ORIGINAL EFFECTIVE DATE: 02/21/19
LAST REVIEW DATE: 02/21/19
LAST CRITERIA REVISION DATE:
ARCHIVE DATE:

TEGSEDI™ (inotersen) subcutaneous injection

Coverage for services, procedures, medical devices and drugs are dependent upon benefit eligibility as outlined in the member's specific benefit plan. This Pharmacy Coverage Guideline must be read in its entirety to determine coverage eligibility, if any.

This Pharmacy Coverage Guideline provides information related to coverage determinations only and does not imply that a service or treatment is clinically appropriate or inappropriate. The provider and the member are responsible for all decisions regarding the appropriateness of care. Providers should provide BCBSAZ complete medical rationale when requesting any exceptions to these guidelines.

The section identified as "Description" defines or describes a service, procedure, medical device or drug and is in no way intended as a statement of medical necessity and/or coverage.

The section identified as "Criteria" defines criteria to determine whether a service, procedure, medical device or drug is considered medically necessary or experimental or investigational.

State or federal mandates, e.g., FEP program, may dictate that any drug, device or biological product approved by the U.S. Food and Drug Administration (FDA) may not be considered experimental or investigational and thus the drug, device or biological product may be assessed only on the basis of medical necessity.

Pharmacy Coverage Guidelines are subject to change as new information becomes available.

For purposes of this Pharmacy Coverage Guideline, the terms "experimental" and "investigational" are considered to be interchangeable.

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

This Pharmacy Coverage Guideline does not apply to FEP or other states' Blues Plans.

Information about medications that require precertification is available at www.azblue.com/pharmacy.

Some large (100+) benefit plan groups may customize certain benefits, including adding or deleting precertification requirements.

All applicable benefit plan provisions apply, e.g., waiting periods, limitations, exclusions, waivers and benefit maximums.

Precertification for medication(s) or product(s) indicated in this guideline requires completion of the request form in its entirety with the chart notes as documentation. All requested data must be provided. Once completed the form must be signed by the prescribing provider and faxed back to BCBSAZ Pharmacy Management at (602) 864-3126 or emailed to Pharmacyprecert@azblue.com. **Incomplete forms or forms without the chart notes will be returned.**

TEGSEDI™ (inotersen) subcutaneous injection (cont.)

Criteria:

- **Criteria for initial therapy:** Tegsedi (inotersen) is considered *medically necessary* and will be approved when **ALL** of the following criteria are met:
1. Prescriber is a Neurologist
 2. Individual is 18 years of age or older
 3. A confirmed diagnosis of polyneuropathy of hereditary transthyretin-mediated amyloidosis
 - Diagnosis confirmed by biopsy and genetic testing documenting pathogenic TTR mutation
 - Signs and symptoms of polyneuropathy
 - Polyneuropathy disability stage \leq 3A
 - Familial amyloid polyneuropathy stage 1 or 2
 4. Individual has failure, contraindication or intolerance to **ALL** the following preferred step therapy agents:
 - Diflunisal 250 mg twice daily used continuously and concurrently with a proton pump inhibitor (lansoprazole, omeprazole, pantoprazole, or rabeprazole) for at least 6 months
 5. **ALL** of the following baseline tests have been completed before initiation of treatment with continued monitoring as clinically appropriate:
 - Platelet count according to REMS
 - Renal function according to REMS
 - Comprehensive metabolic panel
 - Urinary protein to creatinine ratio (UPCR) $<$ 1,000 mg/g
 - eGFR $>$ 45 mL/min/1.73 m²
 6. There are **NO** contraindications
 - Contraindications include:
 - Platelet count $<$ 100 x 10⁹/L
 - History of glomerular nephritis caused by Tegsedi
 - History of hypersensitivity to Tegsedi
 7. Will not be used with Onpattro (patisiran) or tafamidis meglumine
 8. Neuropathy is not due to other causes such as from diabetes mellitus, chronic alcohol, vitamin B12 deficiency, chronic inflammatory demyelinating polyneuropathy
 9. There is vitamin A supplementation using recommended daily allowance
 10. Will not be used in an individual who cannot use a glucocorticoid or immunosuppressants

Initial approval duration: One infusion every week for 6 months

TEGSEDI™ (inotersen) subcutaneous injection (cont.)

- **Criteria for continuation of coverage (renewal request):** Tegsedi (inotersen) is considered *medically necessary* and will be approved when **ALL** of the following criteria are met:
1. Individual continues to be seen by a Neurologist
 2. Individual's condition responded while on therapy
 - Response is defined as:
 - Achieved and maintains at least a 25 % improvement in neurologic function (cranial nerves, reflexes, sensations), motor function (muscle strength), cardiac function (heart rate response to deep breathing, postural blood pressure), quantitative sensory testing (touch-pressure and heat-pain)
 3. Individual has been adherent with the medication
 4. Individual has not developed any contraindications or other significant level 4 adverse drug effects that may exclude continued use
 - Contraindications as listed in the criteria for initial therapy section
 - Significant adverse effect such as:
 - Thrombocytopenia
 - Glomerulonephritis
 - Nephrotic syndrome
 - Stroke
 - Carotid artery dissection
 - Cytokine release symptoms
 - Hepatic dysfunction
 - Detection of treatment emergent anti-platelet IgG antibodies
 5. Will not be used with Onpattro (patisiran) or tafamidis meglumine
 6. There is vitamin A supplementation using recommended daily allowance
 7. Will not be used in an individual who cannot use a glucocorticoid or immunosuppressants

Renewal duration: One infusion every week for 12 months

Description:

Tegsedi (inotersen) is indicated for the treatment of the polyneuropathy of hereditary transthyretin-mediated amyloidosis (hATTR amyloidosis) in adults.

Tegsedi (inotersen) is an antisense oligonucleotide (ASO) that causes degradation of mutant and wild-type TTR mRNA through binding to the TTR mRNA, which results in a reduction of serum TTR protein and TTR protein deposits in tissues. By interfering with the RNA that carries damaged sequences for hATTR amyloidosis, it prevents the formation of amyloid fibrils. Because of the risks of serious bleeding caused by severe thrombocytopenia and because of glomerulonephritis, both of which require frequent monitoring, Patients using

TEGSEDI™ (inotersen) subcutaneous injection (cont.)

Tegsedi (inotersen) will need frequent testing for platelet counts and kidney function before, during and after treatment. Tegsedi (inotersen) is available only through a restricted distribution program under a Risk Evaluation and Mitigation Strategy (REMS) called the TEGSEDI REMS Program.

Background:

Transthyretin amyloidosis is a slowly progressive condition characterized by the buildup of abnormal deposits of amyloid (amyloidosis) in organs and tissues. Protein deposition most frequently occurs in the peripheral nervous system resulting in a loss of sensation in the extremities (peripheral neuropathy). The autonomic nervous system may also be affected by amyloidosis. In some cases, the brain and spinal cord are affected. Other areas of amyloidosis include the heart, kidneys, eyes, and gastrointestinal tract. The age at which symptoms begin to develop varies widely among individuals with this condition, and is typically between ages 20 and 70. The condition is inherited in an autosomal dominant pattern. The disease is caused by a mutation of the *TTR* gene located on chromosome 18 where valine is replaced by methionine at position 30 (*TTR* V30M or Val30Met). There are more than 130 mutations described.

The forms of transthyretin amyloidosis are distinguished symptoms and the body system affected.

The neuropathic form primarily affects the peripheral and autonomic nervous systems, resulting in peripheral neuropathy and difficulty controlling bodily functions. Impairments in bodily functions can include sexual impotence, diarrhea, constipation, problems with urination, and orthostatic hypotension. Some experience heart and kidney problems as well. Various eye problems may occur, such as cloudiness of the clear gel that fills the eyeball (vitreous opacity), dry eyes, glaucoma, or pupils with an irregular or "scalloped" appearance. Some people also develop carpal tunnel syndrome, characterized by numbness, tingling, and weakness in the hands and fingers.

The leptomeningeal form primarily affects the central nervous system. In this form, amyloidosis occurs in the leptomeninges. Protein buildup can cause strokes and hemorrhage, hydrocephalus, ataxia, muscle stiffness and weakness (spastic paralysis), seizures, and loss of intellectual function. Eye problems similar to those in the neuropathic form may also occur.

The cardiac form primarily affects the heart which can lead to arrhythmias, cardiomegaly, or orthostatic hypertension. These abnormalities can lead to progressive heart failure and death. Occasionally, people with the cardiac form of transthyretin amyloidosis have mild peripheral neuropathy.

Various stages of disease have been described. Patients with stage 0 disease are usually asymptomatic but have both a variant form of the *TTR* gene and evidence of amyloid deposits. Patients with stage I (mild) disease are still ambulatory, patients with stage II (moderate) disease are ambulatory but require assistance, and patients with stage III (severe) disease are bedridden or wheelchair bound.

Tissue biopsy should be used to confirm the diagnosis in all cases of amyloidosis, although the diagnosis of amyloidosis may be suspected on the basis of history and clinical manifestations. Tissue biopsy is done using Congo red staining and an immune histochemical study with anti-*TTR* antibodies. Genetic testing is needed to document the pathogenic mutation. If it is normal, *TTR*-FAP is excluded. Current techniques for performing sequence analysis of *TTR*, the only gene known to be associated with *TTR* amyloidosis, detect >99% of disease-causing mutations.

PHARMACY COVERAGE GUIDELINES
SECTION: DRUGS

ORIGINAL EFFECTIVE DATE: 02/21/19
LAST REVIEW DATE: 02/21/19
LAST CRITERIA REVISION DATE:
ARCHIVE DATE:

TEGSEDI™ (inotersen) subcutaneous injection (cont.)

Orthotopic liver transplantation (OLT), which removes the main production site of the amyloidogenic protein, has historically been the standard of care for hereditary TTR amyloidosis. OLT is not effective in the non-neuropathic forms of familial TTR amyloidosis (i.e., cardiac amyloidosis, leptomeningeal amyloidosis, and familial oculoleptomeningeal amyloidosis [FOLMA]).

There are two other modes of treatment. The first is to reduce or halt the amount of mutant transthyretin that is synthesized through gene silencing by the liver. This approach employs use of small interfering RNA (patisiran) and antisense oligonucleotides (inotersen). The other approach is to stabilize mutant tetramers of transthyretin to prevent amyloidogenic monomers. Tetramer stabilizers include diflunisal and tafamidis. Currently under investigation is use of agent(s) to degrade amyloid fibrils that have already been deposited in tissues.

About 3,000 Americans are believed to have polyneuropathy caused by hATTR, which results from abnormally bent and folded proteins produced by mutated RNA. The amyloid fibrils (unusable proteins) deposit in nerves, where they produce pain in the arms, feet, hands and legs. Because they also accumulate in organ tissue, they can enlarge the heart and damage other organ.

Definitions:

Other names:

Familial amyloid polyneuropathy type I (Portuguese-Swedish-Japanese type)
Familial amyloid polyneuropathy type II (Indiana/Swiss or Maryland/German type)
Familial TTR amyloidosis
Amyloid transthyretin polyneuropathy (ATTR-PN)
Familial amyloidotic polyneuropathy (FAP)

Staging:

Clinical staging of TTR-FAP	
Stage 0	No symptoms
Stage I	Unimpaired ambulation; mostly mild sensory, motor, and autonomic neuropathy in the lower limbs
Stage II	Assistance with ambulation required; mostly moderate impairment progression to the lower limbs, upper limbs, and trunk
Stage III	Wheelchair-bound or bedridden; severe sensory, motor, and autonomic involvement of all limbs

Polyneuropathy Disability Staging:

Polyneuropathy Disability Stage	
Stage	Description
0	No symptoms
I	Sensory disturbances but preserved walking capability
II	Impaired walking capacity but ability to walk without a stick or crutches
IIIA	Walking with the help of one stick or crutch
IIIB	Walking with the help of two sticks or crutches
IV	Confined to a wheelchair or bedridden

TEGSEDI™ (inotersen) subcutaneous injection (cont.)

Vitamin A:

Vitamin A (retinol, retinoic acid) is a nutrient important to vision, growth, cell division, reproduction and immunity. Vitamin A is found in many foods, such as spinach, dairy products and liver. Other sources are foods rich in beta-carotene, such as green leafy vegetables, carrots and cantaloupe. Your body converts beta-carotene into vitamin A. The recommended daily amount of vitamin A is 900 micrograms (mcg) for adult men and 700 mcg for adult women.

<https://ods.od.nih.gov/factsheets/VitaminA-Consumer/>

Currently, vitamin A is listed on food and supplement labels in international units (IUs) even though nutrition scientists rarely use this measure. Conversion rates between mcg retinol activity equivalents (RAE) and IU are as follows:

- 1 IU retinol = 0.3 mcg RAE
- 1 IU beta-carotene from dietary supplements = 0.15 mcg RAE
- 1 IU beta-carotene from food = 0.05 mcg RAE
- 1 IU alpha-carotene or beta-cryptoxanthin = 0.025 mcg RAE

Resources:

Onpattro (patisiran) product information accessed 11-13-18 at DailyMed:

<https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=e87ec36f-b4b4-49d4-aea4-d4ffb09b0970>

Tedsedi (inotersen) product information accessed 11-13-18 at DailyMed:

<https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=8513207e-b55f-417b-9473-af785146a543>

UpToDate: Overview of amyloidosis. Current through Oct 2018. https://www.uptodate.com.mwu.idm.oclc.org/contents/overview-of-amyloidosis?search=hereditary%20transthyretin-mediated%20amyloidosis&source=search_result&selectedTitle=2~150&usage_type=default&display_rank=2

UpToDate: Genetic factors in the amyloid diseases. Current through Oct 2018. https://www.uptodate.com.mwu.idm.oclc.org/contents/genetic-factors-in-the-amyloid-diseases?topicRef=5589&source=see_link

UpToDate: Treatment of amyloid cardiomyopathy. Current through Oct 2018. https://www.uptodate.com.mwu.idm.oclc.org/contents/treatment-of-amyloid-cardiomyopathy?sectionName=Specific%20therapy%20of%20TTR%20amyloidosis&topicRef=5589&anchor=H194831834&source=see_link#H1994831834

Berk JL, Suhr OB, Obici L, et al.: Repurposing Diflunisal for Familial Amyloid Polyneuropathy A Randomized Clinical Trial. JAMA 2013; 310 (24):2658-2667.

Ando Y, Coelho T, Berk JL, et al.: Guideline of transthyretin-related hereditary amyloidosis for clinicians. Orphanet Journal of Rare Diseases 2013; 8:31.

Sekijima Y. Recent progress in the understanding and treatment of transthyretin amyloidosis. J Clinical Pharmacy and Therapeutics, 2014; 39: 225–233.



PHARMACY COVERAGE GUIDELINES
SECTION: DRUGS

ORIGINAL EFFECTIVE DATE: 02/21/19
LAST REVIEW DATE: 02/21/19
LAST CRITERIA REVISION DATE:
ARCHIVE DATE:

TEGSEDI™ (inotersen) subcutaneous injection (cont.)

Hawkins PN , Ando Y, Dispenzeri A, et al.: Evolving landscape in the management of transthyretin amyloidosis. *Annals of Medicine* 2015; 47: 625–638.

Sekijima Y, Ueda M, Koike H, et al.: Diagnosis and management of transthyretin familial amyloid polyneuropathy in Japan: red-flag symptom clusters and treatment algorithm. *Orphanet Journal of Rare Diseases* 2018; 13:6.

Adams D, Gonzalez-Duarte A, O’Riordan WD, et al.: Patisiran, an RNAi Therapeutic, for Hereditary Transthyretin Amyloidosis. *NEJM* 2018; 379:11-21.

Benson MD, Waddington-Cruz M, Berk JL, et al: Inotersen Treatment for Patients with Hereditary Transthyretin Amyloidosis. *NEJM* 2018; 379:22-31.

Maurer MS, Schwartz JH, Gundapaneni B, et al.: Tafamidis Treatment for Patients with Transthyretin Amyloid Cardiomyopathy. *NEJM* 2018; 379:1007-1016.



An Independent Licensee of the Blue Cross and Blue Shield Association

Fax completed prior authorization request form to 602-864-3126 or email to pharmacyprecert@azblue.com.
 Call 866-325-1794 to check the status of a request.
 All requested data must be provided. **Incomplete forms or forms without the chart notes will be returned.**
 Pharmacy Coverage Guidelines are available at www.azblue.com/pharmacy.

Pharmacy Prior Authorization Request Form

Do not copy for future use. Forms are updated frequently.

REQUIRED: Office notes, labs, and medical testing relevant to the request that show medical justification are required.

Member Information			
Member Name (first & last):	Date of Birth:	Gender:	BCBSAZ ID#:
Address:	City:	State:	Zip Code:
Prescribing Provider Information			
Provider Name (first & last):	Specialty:	NPI#:	DEA#:
Office Address:	City:	State:	Zip Code:
Office Contact:	Office Phone:	Office Fax:	
Dispensing Pharmacy Information			
Pharmacy Name:	Pharmacy Phone:	Pharmacy Fax:	
Requested Medication Information			
Medication Name:	Strength:	Dosage Form:	
Directions for Use:	Quantity:	Refills:	Duration of Therapy/Use:
<input type="checkbox"/> Check if requesting brand only <input type="checkbox"/> Check if requesting generic			
<input type="checkbox"/> Check if requesting continuation of therapy (prior authorization approved by BCBSAZ expired)			
Turn-Around Time For Review			
<input type="checkbox"/> Standard <input type="checkbox"/> Urgent. Sign here: _____ <input type="checkbox"/> Exigent (requires prescriber to include a written statement)			
Clinical Information			
1. What is the diagnosis? Please specify below. ICD-10 Code: _____ Diagnosis Description: _____			
2. <input type="checkbox"/> Yes <input type="checkbox"/> No Was this medication started on a recent hospital discharge or emergency room visit?			
3. <input type="checkbox"/> Yes <input type="checkbox"/> No There is absence of ALL contraindications.			
4. What medication(s) has the individual tried and failed for this diagnosis? Please specify below. Important note: Samples provided by the provider are not accepted as continuation of therapy or as an adequate trial and failure.			
Medication Name, Strength, Frequency	Dates started and stopped or Approximate Duration	Describe response, reason for failure, or allergy	
5. Are there any supporting labs or test results? Please specify below.			
Date	Test	Value	

Pharmacy Prior Authorization Request Form

6. Is there any additional information the prescribing provider feels is important to this review? Please specify below.
For example, explain the negative impact on medical condition, safety issue, reason formulary agent is not suitable to a specific medical condition, expected adverse clinical outcome from use of formulary agent, or reason different dosage form or dose is needed.

Signature affirms that information given on this form is true and accurate and reflects office notes

Prescribing Provider's Signature: _____ Date: _____

Please note: Some medications may require completion of a drug-specific request form.

Incomplete forms or forms without the chart notes will be returned.

Office notes, labs, and medical testing relevant to the request that show medical justification are required.