



An Independent Licensee of the Blue Cross Blue Shield Association

PHARMACY COVERAGE GUIDELINES
SECTION: DRUGS

ORIGINAL EFFECTIVE DATE: 1/22/2015
LAST REVIEW DATE: 2/17/2022
LAST CRITERIA REVISION DATE: 2/17/2022
ARCHIVE DATE:

GLEEVEC® (imatinib mesylate) oral tablet Imatinib Mesylate oral tablet

Coverage for services, procedures, medical devices and drugs are dependent upon benefit eligibility as outlined in the member's specific benefit plan. This Pharmacy Coverage Guideline must be read in its entirety to determine coverage eligibility, if any.

This Pharmacy Coverage Guideline provides information related to coverage determinations only and does not imply that a service or treatment is clinically appropriate or inappropriate. The provider and the member are responsible for all decisions regarding the appropriateness of care. Providers should provide BCBSAZ complete medical rationale when requesting any exceptions to these guidelines.

The section identified as "Description" defines or describes a service, procedure, medical device or drug and is in no way intended as a statement of medical necessity and/or coverage.

The section identified as "Criteria" defines criteria to determine whether a service, procedure, medical device or drug is considered medically necessary or experimental or investigational.

State or federal mandates, e.g., FEP program, may dictate that any drug, device or biological product approved by the U.S. Food and Drug Administration (FDA) may not be considered experimental or investigational and thus the drug, device or biological product may be assessed only on the basis of medical necessity.

Pharmacy Coverage Guidelines are subject to change as new information becomes available.

For purposes of this Pharmacy Coverage Guideline, the terms "experimental" and "investigational" are considered to be interchangeable.

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

This Pharmacy Coverage Guideline does not apply to FEP or other states' Blues Plans.

Information about medications that require precertification is available at www.azblue.com/pharmacy.

Some large (100+) benefit plan groups may customize certain benefits, including adding or deleting precertification requirements.

All applicable benefit plan provisions apply, e.g., waiting periods, limitations, exclusions, waivers and benefit maximums.

Precertification for medication(s) or product(s) indicated in this guideline requires completion of the [request form](#) in its entirety with the chart notes as documentation. **All requested data must be provided.** Once completed the form must be signed by the prescribing provider and faxed back to BCBSAZ Pharmacy Management at (602) 864-3126 or emailed to Pharmacyprecert@azblue.com. **Incomplete forms or forms without the chart notes will be returned.**



An Independent Licensee of the Blue Cross Blue Shield Association

PHARMACY COVERAGE GUIDELINES
SECTION: DRUGS

ORIGINAL EFFECTIVE DATE: 1/22/2015
LAST REVIEW DATE: 2/17/2022
LAST CRITERIA REVISION DATE: 2/17/2022
ARCHIVE DATE:

GLEEVEC® (imatinib mesylate) oral tablet Imatinib Mesylate oral tablet

Criteria:

- **Criteria for initial therapy:** Gleevec (imatinib mesylate) or Imatinib mesylate is considered **medically necessary** and will be approved when **ALL** of the following criteria are met:
1. Prescriber is a physician specializing in the patient's diagnosis or is in consultation with an Oncologist, HIV/AIDS Specialist, or Gastroenterologist depending upon indication or use
 2. A diagnosis of **ONE** of the following:
 - a. Newly diagnosed adult and pediatric (1 year of age or older) patient with Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in chronic phase
 - b. Philadelphia chromosome positive chronic myeloid leukemia in blast crisis, accelerated phase, or in chronic phase after failure of interferon-alpha therapy
 - c. Adult (18 years of age or older) patient with relapsed or refractory Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL)
 - d. Pediatric (1 year of age or older) patient with newly diagnosed Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL) in combination with chemotherapy
 - e. Adult patient (18 years of age or older) with myelodysplastic/myeloproliferative diseases associated with platelet-derived growth factor receptor (PDGFR) gene re-arrangements
 - f. Adult patient (18 years of age or older) with aggressive systemic mastocytosis without the D816V c-Kit mutation or with c-Kit mutational status unknown
 - g. Adult patient (18 years of age or older) with hypereosinophilic syndrome and/or chronic eosinophilic leukemia (HES/CEL) who have the FIP1L1-PDGFR α fusion kinase (mutational analysis or fluorescence in situ hybridization [FISH] demonstration of CHIC2 allele deletion) and for patients with HES and/or CEL who are FIP1L1-PDGFR α fusion kinase negative or unknown
 - h. Adult patient (18 years of age or older) with unresectable, recurrent and/or metastatic dermatofibrosarcoma protuberans (DFSP)
 - i. Adult patient (18 years of age or older) with Kit (CD117) positive unresectable and/or metastatic malignant gastrointestinal stromal tumors (GIST)
 - j. Adjuvant treatment of adult patient (18 years of age or older) following resection of Kit (CD117) positive GIST
 - k. Other request for a specific oncologic direct treatment use that is found and listed in the National Comprehensive Cancer Network (NCCN) Guidelines with Categories of Evidence and Consensus of 1 and 2A



An Independent Licensee of the Blue Cross Blue Shield Association

PHARMACY COVERAGE GUIDELINES
SECTION: DRUGS

ORIGINAL EFFECTIVE DATE: 1/22/2015
LAST REVIEW DATE: 2/17/2022
LAST CRITERIA REVISION DATE: 2/17/2022
ARCHIVE DATE:

GLEEVEC® (imatinib mesylate) oral tablet Imatinib Mesylate oral tablet

3. **ALL** of the following baseline tests have been completed before initiation of treatment:
 - a. Where applicable, genetic testing has been completed using an FDA approved test and the result of testing is submitted
 - b. Other required testing as outlined by manufacturer and FDA labeling have been completed and/or are ongoing
 - c. Liver function tests
 - d. Assessment of hydration status and uric acid levels, with correction if abnormal
 - e. Negative pregnancy test in a woman of childbearing age
4. Request for **brand** Gleevec: individual has failure, contraindication or intolerance to **generic imatinib mesylate**
5. There are no significant interacting drugs

Initial approval duration: 6 months

- **Criteria for continuation of coverage (renewal request):** Gleevec (imatinib mesylate) or Imatinib mesylate is considered **medically necessary** and will be approved when **ALL** of the following criteria are met:
1. Individual continues to be seen by a physician specializing in the patient's diagnosis or is in consultation with an Oncologist, HIV/AIDS Specialist, or Gastroenterologist depending upon indication or use
 2. Individual's condition has responded while on therapy
 - a. Response is defined as:
 - i. Documented evidence of efficacy, disease stability and/or improvement
 - ii. No evidence individual has developed any significant unacceptable adverse drug reactions that may exclude continued use
 3. Individual has been adherent with the medication
 4. Individual has not developed any significant adverse drug effects that may exclude continued use
 - a. Significant adverse effect such as:
 - i. Cytopenias (anemia, neutropenia, thrombocytopenia)
 - ii. Pleural effusions, pericardial effusions, pulmonary edema, ascites:
 - iii. Heart failure, left ventricular dysfunction, or cardiogenic shock
 - iv. Hepatotoxicity
 - v. GI bleeding or perforation
 - vi. Erythema multiforme/Stevens-Johnson Syndrome
 - vii. Tumor lysis syndrome
 - viii. Renal toxicity



An Independent Licensee of the Blue Cross Blue Shield Association

PHARMACY COVERAGE GUIDELINES
SECTION: DRUGS

ORIGINAL EFFECTIVE DATE: 1/22/2015
LAST REVIEW DATE: 2/17/2022
LAST CRITERIA REVISION DATE: 2/17/2022
ARCHIVE DATE:

GLEEVEC® (imatinib mesylate) oral tablet Imatinib Mesylate oral tablet

5. There are no significant interacting drugs

Renewal duration: 12 months

➤ Criteria for a request for non-FDA use or indication, treatment with dosing, frequency, or duration outside the FDA-approved dosing, frequency, and duration, refer to one of the following Pharmacy Coverage Guideline:

1. **Off-Label Use of Non-cancer Medications**
 2. **Off-Label Use of Cancer Medications**
-

Description:

Gleevec® (imatinib) is used for the treatment of several malignancies: acute lymphoblastic leukemia, aggressive systemic mastocytosis, chronic myeloid leukemia, dermatofibrosarcoma protuberans, gastrointestinal stromal tumors, hypereosinophilic syndrome / chronic eosinophilic leukemia, and myelodysplastic / myeloproliferative disease. It is a small molecule tyrosine kinase inhibitor with several important actions on cellular function. It blocks tyrosine kinase activity of several key proteins involved the regulation of growth, differentiation, and apoptosis. Deregulation of tyrosine kinase activity has been shown to play an important role in development of various cancers.

Tyrosine kinase inhibitors (TKIs) are a class of agents designed to compete with adenosine triphosphate (ATP) for its binding pocket within the intracellular domain of wild type and/or mutated receptor. Binding of Imatinib within the pocket blocks downstream signaling important for tumor growth. All TKIs are designed to compete with ATP for the ATP binding pocket of similar or different tyrosine kinases that are mutated and/or over-expressed in specific tumors.

In the treatment of chronic myeloid leukemia (CML), Imatinib inhibits the breakpoint cluster region-Abelson (BCR-ABL) tyrosine kinase fusion protein created by the chromosomal abnormality known as the Philadelphia chromosome (Ph). BCR-ABL is uniquely expressed by leukemic cells and is essential for the survival of these cells. The fusion protein is present in 95% of individuals with CML. Philadelphia chromosome is also an abnormality seen in approximately 30% of newly diagnosed adults with acute lymphoblastic leukemia (ALL). Imatinib potently and specifically inhibits growth of BCR-ABL expressing cells leading to inhibition of proliferation and apoptosis in BCR-ABL positive cell lines as well as fresh leukemic cells.

Gastrointestinal stromal tumors (GISTs) are neoplasms of the gastrointestinal (GI) tract. They are thought to arise from the interstitial cells of Cajal. GISTs are defined by the expression of the tyrosine kinase c-KIT (CD117) receptor, the receptor for stem cell factor (SCF), in the tumor cells resulting in constitutive activation of the tyrosine kinase. The c-KIT is expressed in approximately 85% of GISTs. Imatinib inhibits proliferation and induces apoptosis in GISTs cells, which express an activating c-KIT mutation.

PHARMACY COVERAGE GUIDELINES
SECTION: DRUGS

ORIGINAL EFFECTIVE DATE: 1/22/2015
LAST REVIEW DATE: 2/17/2022
LAST CRITERIA REVISION DATE: 2/17/2022
ARCHIVE DATE:

GLEEVEC® (imatinib mesylate) oral tablet **Imatinib Mesylate oral tablet**

Mutation of c-KIT is also found in the myeloproliferative disorder systemic mastocytosis. In GISTs, mutations and deletions of c-KIT are typically found in the juxta membrane domain, resulting in constitutive activation of the tyrosine kinase. With systemic mastocytosis, the characteristic D816V activating c-KIT mutation is within the kinase domain itself. While Imatinib has significant activity in advanced GISTs, it has proven largely unsuccessful in the treatment of systemic mastocytosis due to ineffective targeting of c-KIT kinases with the D816V mutation. All responses in patients with systemic mastocytosis were seen in those who were negative for D816V c-KIT mutation.

The idiopathic hypereosinophilic syndrome (HES), now reclassified as chronic eosinophilic leukemia (CEL), is characterized by the expression of the FIP1-like-1–platelet-derived growth factor receptor alpha (FIP1L1-PDGFR α) fusion protein, which is generated by an interstitial chromosomal deletion and results in constitutive signaling through PDGFR α . Dermatofibrosarcoma protuberans (DFSP) is a rare soft tissue tumor characterized by the presence of a distinctive, reciprocal rearrangement of certain chromosomes. The rearrangement leads to the fusion of collagen type 1 alpha-1 (COL1A1) chain to platelet-derived growth factor beta (PDGFB). The formation of COL1A1-PDGFB fusion gene results in constitutional up-regulation of PDGFB expression, leading to continuous autocrine activation of the receptor. Imatinib is an inhibitor specific for platelet derived growth factor receptor and is effect for HES/CEL and DFSP.

Definitions:

BCR-ABL1 (IS) Response Milestones:

BCR-ABL1 (IS)	3 months	6 months	12 months	> 12 months
> 10%	YELLOW	RED		
>1-10%	GREEN		YELLOW	RED
>0.1-1%	GREEN			YELLOW
≤ 0.1%	GREEN			
	Clinical considerations		2 nd line & subsequent treatment options	
Red	<ul style="list-style-type: none"> Evaluate compliance & drug interactions Mutational analysis 		<ul style="list-style-type: none"> Switch to alternate TKI Evaluate for HCT 	
Yellow	<ul style="list-style-type: none"> Evaluate compliance & drug interactions Mutational analysis 		<ul style="list-style-type: none"> Switch to alternate TKI or continue same TKI or dose escalation of imatinib (to max of 800 mg) Evaluate for HCT 	
Green	<ul style="list-style-type: none"> Monitor response & side effects 		<ul style="list-style-type: none"> Continue same TKI 	

Accelerated Phase CML:

Modified Criteria used at MD Anderson Cancer Center (most commonly used in clinical trials)
Peripheral blood blasts \geq 15% and < 30%
Peripheral blood blasts and promyelocytes combined \geq 30%
Peripheral blood basophils \geq 20%
Platelet count \leq 100 x 10 ⁹ /L unrelated to therapy
Additional clonal cytogenetic abnormalities in Ph+ cells
Semin Hematol 1988;25:49-61

PHARMACY COVERAGE GUIDELINES
SECTION: DRUGS

ORIGINAL EFFECTIVE DATE: 1/22/2015
LAST REVIEW DATE: 2/17/2022
LAST CRITERIA REVISION DATE: 2/17/2022
ARCHIVE DATE:

GLEEVEC® (imatinib mesylate) oral tablet

Imatinib Mesylate oral tablet

Br J Haematol 1997;99:30-35
 Blood 1993;82:691-703
 Blood 2002;99:1928-1937

Blast Phase CML:

World Health Organization Criteria	International Bone Marrow Transplant Registry
Blasts \geq 20% of peripheral white blood cells or of nucleated bone marrow cells Extramedullary blast proliferation Large foci or clusters of blasts in the bone marrow biopsy	\geq 30% blasts in the blood, marrow, or both Extramedullary infiltrates or leukemic cells
NCCN Chronic myeloid leukemia. Version 1.2018, July 26, 2017	

Treatment options based on BCR-ABL1 mutation profile:

Mutation	Treatment recommendations
E255K/V, F359V/C/I or Y253H	Dasatinib
F317L/V/I/C, T315A, or V299L	Nilotinib
E255K/V, F317L/V/I/C, F359V/C/I, T315A, or Y253H	Bosutinib
T315I	Ponatinib, Omacetaxine, allogeneic HCT, or clinical trial

- Patients with disease that is resistant to primary treatment with imatinib should be treated with nilotinib, dasatinib, or bosutinib in the second-line setting.
- Patients with disease that is resistant to primary treatment with nilotinib or dasatinib could be treated with an alternative TKI (other than imatinib) in the second-line setting.
- Ponatinib is also a treatment option for patients for whom no other TKI is indicated.
- Omacetaxine is a treatment option for patients with disease that is resistant and/or intolerant to 2 or more TKIs.

Definitions for response and relapse in CML:

CHR	Complete normalization of peripheral blood counts with leukocyte count $<$ $10 \times 10^9/L$ Platelet count $<$ $450 \times 10^9/L$ No immature cells (such as myelocytes, promyelocytes, or blasts) in peripheral blood No signs & symptoms of disease, with disappearance of palpable splenomegaly
CyR	Complete CyR (CCyR): no Ph+ metaphases (correlates to <i>BCR-ABL</i> (IS) \leq 1% ($>$ 0.1-1%)) Partial CyR (PCyR): 1-35% Ph+ metaphases Major CyR: 0-35% Ph+ metaphases Minor CyR: $>$ 35% Ph+ metaphases No response: $>$ 95% Ph+ metaphases
MR	Early MR (EMR) – <i>BCR-ABL</i> (IS) \leq 10% at 3 and 6 months Major MR (MMR) – <i>BCR-ABL</i> (IS) \leq 0.1% or \geq 3 log reduction in <i>BCR-ABL1</i> mRNA from the standardized baseline, if qPCR (IS) is not available Complete MR (CMR) – is variably described, and is best defined by the assay's level of sensitivity (such as MR 4.5)
Relapse	Any sign of loss of response define as hematologic or cytogenetic 1 log increase in <i>BCR-ABL1</i> transcript levels with loss of MMR should prompt bone marrow evaluation for loss of CCyR but is not itself defined as relapse (hematologic or cytogenetic relapse)

CHR: complete hematologic response
 CyR: cytogenetic response
 MR: molecular response
 IS: International scale – the ratio of the BCR-ABL1 transcriptions to ABL1 transcriptions

GLEEVEC® (imatinib mesylate) oral tablet **Imatinib Mesylate oral tablet**

Molecular response International Scale:

International Scale (IS)	
MR 2	Detectable disease at a level of $\leq 1\%$ on the IS (≥ 2 log reduction from the standardized baseline). This level of response roughly corresponds to a "complete cytogenetic response"
MR 3	Detectable disease at a level of $\leq 0.1\%$ on the IS (≥ 3 log reduction from the standardized baseline). This level of response has been termed a "major molecular response"
MR 4	Either detectable disease at a level of $\leq 0.01\%$ on the IS (≥ 4 log reduction) or undetectable disease in cDNA with $\geq 10,000$ ABL1 transcripts. This level of response requires that the assay being used is sensitive enough to detect a single abnormal transcript amongst 10,000 normal ABL1 transcripts
MR 4.5	Either detectable disease at a level of $\leq 0.0032\%$ on the IS (≥ 4.4 log reduction) or undetectable disease in cDNA with $\geq 32,000$ ABL1 transcripts. This level of response requires that the assay being used is sensitive enough to detect a single abnormal transcript amongst 32,000 normal ABL1 transcripts

Monitoring Response to TKI Therapy and Mutational Analysis:

Test	Recommendation
Bone marrow cytogenetic	<ul style="list-style-type: none"> • At diagnosis • Failure to reach response milestone • Any signs of loss of response (defined as hematologic or cytogenetic relapse)
Quantitative RT-PCT (qPCR) using IS	<ul style="list-style-type: none"> • At diagnosis • Every 3 months after initiating treatment. After <i>BCR-ABL1</i> (IS) $\leq 1\%$ ($> 0.1-1\%$) has been achieved, every 3 months x 2 y and every 3-6 months thereafter • If there is a 1-log increase in <i>BCR-ABL1</i> transcript levels with MMR, qPCR should be repeated in 1-3 months
<i>BCR-ABL1</i> kinase domain mutation analysis	<ul style="list-style-type: none"> • Chronic phase <ul style="list-style-type: none"> ➢ Failure to reach response milestone ➢ Any signs of loss of response (defined as hematologic or cytogenetic relapse) ➢ 1-log increase in <i>BCR-ABL1</i> transcript levels and loss of MMR • Disease progression to accelerated or blast phase

Resources:

Gleevec (imatinib mesylate) product information, revised by Novartis Pharmaceuticals Corporation 08-2020. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed December 11, 2021.

Imatinib mesylate product information, revised by Lupin Pharmaceuticals, Inc. 10-2020. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed December 11, 2021.

National Comprehensive Cancer Network (NCCN) Clinical Practice Guidelines in Oncology (NCCN Guidelines®): Acute Lymphoblastic Leukemia Version 2.2021 – Updated July 19, 2021. Available at <https://www.nccn.org>. Accessed December 11, 2021.

National Comprehensive Cancer Network (NCCN) Clinical Practice Guidelines in Oncology (NCCN Guidelines®): Pediatric Acute Lymphoblastic Leukemia Version 1.2022 – Updated October 01, 2021. Available at <https://www.nccn.org>. Accessed December 11, 2021.



An Independent Licensee of the Blue Cross Blue Shield Association

PHARMACY COVERAGE GUIDELINES
SECTION: DRUGS

ORIGINAL EFFECTIVE DATE: 1/22/2015
LAST REVIEW DATE: 2/17/2022
LAST CRITERIA REVISION DATE: 2/17/2022
ARCHIVE DATE:

GLEEVEC® (imatinib mesylate) oral tablet Imatinib Mesylate oral tablet

National Comprehensive Cancer Network (NCCN) Clinical Practice Guidelines in Oncology (NCCN Guidelines®): Chronic Myeloid Leukemia Version 2.2022 – Updated November 15, 2021. Available at <https://www.nccn.org>. Accessed December 11, 2021.

National Comprehensive Cancer Network (NCCN) Clinical Practice Guidelines in Oncology (NCCN Guidelines®): Dermatofibrosarcoma Protuberans Version 1.2022 – Updated November 17, 2021. Available at <https://www.nccn.org>. Accessed December 11, 2021.

National Comprehensive Cancer Network (NCCN) Clinical Practice Guidelines in Oncology (NCCN Guidelines®): Gastrointestinal Stromal Tumors Version 1.2021 – Updated October 30, 2020. Available at <https://www.nccn.org>. Accessed December 11, 2021.

National Comprehensive Cancer Network (NCCN) Clinical Practice Guidelines in Oncology (NCCN Guidelines®): Myelodysplastic Syndromes Version 2.2022 – Updated November 15, 2021. Available at <https://www.nccn.org>. Accessed December 11, 2021.

National Comprehensive Cancer Network (NCCN) Clinical Practice Guidelines in Oncology (NCCN Guidelines®): Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Fusion Genes Version 4.2021 – Updated July 09, 2021. Available at <https://www.nccn.org>. Accessed December 11, 2021.

National Comprehensive Cancer Network (NCCN) Clinical Practice Guidelines in Oncology (NCCN Guidelines®): Systemic Mastocytosis Version 3.2021 – Updated July 09, 2021. Available at <https://www.nccn.org>. Accessed December 11, 2021.

Off Label Use of Cancer Medications: A.R.S. §§ 20-826(R) & (S). Subscription contracts; definitions.

Off Label Use of Cancer Medications: A.R.S. §§ 20-1057(V) & (W). Evidence of coverage by health care service organizations; renewability; definitions.